




Egyptian Society of Radiology and Nuclear Medicine
The Egyptian Journal of Radiology and Nuclear Medicine

www.elsevier.com/locate/ejrnrm
www.sciencedirect.com



ORIGINAL ARTICLE

Comparative study between black blood T2* and conventional bright GRE sequences in assessment of myocardial iron concentration



Nivan Hany Khater^a, Hazem Mohamed El Shahat^b, Hadeer Safwat Fahmy^{b,*},
Samah Saleh El Hadidy^c

^a Ain Shams University Hospitals, Cairo, Egypt

^b Zagazig University Hospitals, Radiology department, Zagazig, Egypt

^c Mansoura University Hospital, Community Medicine Department, Mansoura, Egypt

Received 19 November 2015; accepted 7 January 2016

Available online 3 February 2016

KEYWORDS

Black blood;
Myocardial;
Iron;
Concentration

Abstract *Purpose:* To compare the efficacy and reproducibility of black blood T2* to the conventional bright blood sequence in the assessment of myocardial iron concentration.

Materials and methods: We performed both conventional bright blood and black blood MRI T2* sequences in 50 thalassemia major patients, the results were statistically analyzed to assess the correlation of techniques, study reproducibility and interobserver agreement.

Results: Cardiac T2* values ranged from 2.39 to 47.9 ms using bright blood sequence and 2.07 to 46.81 using the black blood sequence. There was positive significant correlation of both sequences in all patients. However the black blood technique was superior to bright blood technique as regards the study reproducibility ($R^2 1.9 \pm$ versus 2.4 ± 14.7 $p < 0.001$) in addition to the better inter-observer agreement of black blood technique compared to the bright blood technique (3.2 ± 1.2 versus 8.3 ± 2.4 $p < 0.001$).

Conclusions: Black blood T2* technique provides clearly defined septal borders, avoids bright blood signal contamination of the myocardium, has superior study reproducibility and inter-observer agreement which favors this technique in the assessment of iron myocardial concentration.

© 2016 The Egyptian Society of Radiology and Nuclear Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Patients of thalassemia major on regular blood transfusions suffer from inevitable accumulation of iron in tissues of the heart, liver, and endocrine systems. Myocardial iron deposition resulting in cardiac failure and death at a young age. Early detection and assessment of myocardial iron is crucial to

* Corresponding author.

E-mail addresses: Nivan_khater@yahoo.com (N.H. Khater), Hazem_el_shahat@hotmail.com (H.M. El Shahat), Hadeer_fahmy68@hotmail.com (H.S. Fahmy), Samah.elhadidy@yahoo.com (S.S. El Hadidy).

Peer review under responsibility of The Egyptian Society of Radiology and Nuclear Medicine.

<http://dx.doi.org/10.1016/j.ejrnrm.2016.01.006>

0378-603X © 2016 The Egyptian Society of Radiology and Nuclear Medicine. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

evaluate the risk of cardiac complications and tailoring appropriate iron chelating treatment (1).

Despite the fact that transfusions improve health and survival, however the consequent tissue iron deposition causes organ damage in the long term. Patients treated with regular blood transfusions usually die early from cardiac failure secondary to myocardial siderosis (2).

Cardiomyopathy can be treated if intensive iron chelation treatment is initiated properly (3), but diagnosis is often delayed by the concealed cardiac iron deposition and the late development of symptoms, and echocardiographic abnormalities (4). Once heart failure develops, the prognosis is usually poor with abrupt deterioration and death, even with intensive chelation therapy (5).

Early diagnosis and measurement of myocardial iron deposition would allow proper management and help to reduce mortality from this reversible cardiomyopathy; however, endomyocardial biopsy is inaccurate for measuring myocardial iron because of high rate of sampling error with very small biopsy samples (6,15).

The need for an alternative noninvasive measurement of myocardial iron led to the development of an optimized cardiac T_2^* magnetic resonance (MR) technique (7).

Intracellular iron particles deposition causes shortening of the MR relaxation parameter T_2^* with increase in its reciprocal, R_2^* due to microscopic magnetic field in-homogeneity.

T_2^* decay is measured by gradient echo CMR, and iron overload is diagnosed when the values are less than 20 ms (8). Recently, single breath-hold sequence was evolved with shorter scan time, better image registration between images, and excellent reported reproducibility (9). This technique has become the standard for myocardial iron concentration assessment and has also proved value in the assessment of chelation regimens (10).

The white blood technique, however, has some undesirable characteristics including poor contrast between blood pool and myocardial margins, motion artifacts and blood flow compromising measurement accuracy. Recently, a black blood' double inversion recovery sequence which suppresses the blood signal, has reported a preliminary good reproducibility compared to the white blood T_2^* imaging technique (11).

2. Objectives

Our study aimed to compare the efficacy and the technical reproducibility of black blood T_2^* and the conventional bright blood sequence in the assessment of myocardial iron concentration.

3. Patients and methods

A prospective two-year study of 50 patients on regular blood transfusion, referred for MRI assessment of cardiac siderosis has undergone an imaging protocol which includes measurement of cardiac T_2^* using pulse sequences without and with a blood suppression prepulse to evaluate interstudy reproducibility.

All the patients were known TM and were on regular blood transfusion for more than 18 months. Between the age range 8 and 41 years (mean 27) 56% were male.

17 patients (34%) showed cardiac siderosis ($T_2^* < 20$ ms).

Our patients were examined by a 1.5 T scanner (Siemens Espree, Erlangen, Germany) with a 8 channel phased array cardiac coil and ECG gating.

Three mid-ventricular short axis scans with 6 mm slice thickness were obtained. For bright blood acquisition, we used a multi echo gradient sequence (matrix 128×256 pixels, field of view 40 cm, and flip angle 20°). The short axis images were taken with a single breath-hold (8–12 s) at 8 different echo times ranged from 2.54 to 17.90 ms at 2 ms increments. Images of white blood technique were scanned immediately after the R-wave, while for black blood acquisition, we apply a double inversion recovery pulse on the R-wave with prolonged inversion time to diastole, that developing 8 increasing echo time sets of images (12).

We analyze our images using Thalassemia CMR tools (Cardiovascular Imaging Solutions, London, UK). Our ROI full thickness section included the ventricular septum but avoiding the blood pool cavity and cardiac vessels (13,14).

T_2^* was acquired using an exponential curve fit, $SI = SI_0 \cdot \exp(TE/T_2^*)$, and the back ground noise in delayed echo times was detected in bright blood technique only, and was removed using the truncation method for better curve fit. The image quality and the presence of artifacts were assessed in all sequences.

3.1. Statistical methods

Data were statistically analyzed using SPSS (Statistical Package for Social Sciences version 20.0). First, the excel file was imported then full labeling of all variables and their values were performed with exploration of data for any odd or abnormal values before performing the preliminary analysis.

Line graph was utilized to present the correlation between bright blood and black blood techniques.

We apply Kaba test in the analysis and the assessment of image quality and a p value of <0.05 was considered significant.

4. Results

All 50 patients included in this study had thalassemia major anemia and were receiving regular blood transfusion. Between the age range 8 and 41, means $SD 23.38 \pm 8.8$, 28 were males and 22 females.

Table 1 shows characteristics of the patients ($N = 50$), showing 56% of the patients aged from 20 to 39 years with Mean \pm SD (23.38 ± 8.8 years). Most of the patients were males.

Table 1 Age and sex distribution of 50 patients.

Characteristics	Patient	
	$N = 50$	%
<i>Gender</i>		
Male	28	56
Female	22	44
Mean \pm ST. deviation		
<i>Age</i>		
Range	8–41	
Mean \pm ST. deviation	23.38 \pm 8.8	

4.1. T2* myocardial iron concentration results of black blood and bright conventional sequences

Out of 50 patients 33 were normal, while 17 showed varying degree of myocardial iron concentration: 6 were light, 6 moderate while 5 showed severe iron myocardial concentration. Table 2 shows that the majority of our patients (33%) were non cardiac siderosis and only (17%) had severe degree of iron deposition.

The values of cardiac T2* ranged from 2.39–47.9 ms as measured with bright blood acquisition and 2.07–46.81 ms using the black blood sequence.

We reported no significant difference between the mean cardiac T2* values for all 50 patients using both the black blood and the conventional bright blood sequences (30.46 ms versus 30.12 ms; $p = 0.26$).

Table 2 Distribution of disease severity among study population.

Grade (severity)	Patients	
	$N = 50$	%
Normal	33	66
Light	6	12
Moderate	6	12
Severe	5	10

Table 3 Cardiac T2* values measured by bright and black blood techniques.

Myocardial iron concentration (g/ms)	Bright blood	Black blood
Normal	29.59–47.95	30.22–46.81
Light	16.30–18.72	16.48–18.11
Moderate	12.45–13.88	12.04–12.67
Severe	2.39–4.56	2.07–4.21
Mean \pm SD	30.12 \pm 14.7	30.46 \pm 1.47
R	0.99	
p value	0.000 ^a	

^a High significance.

Table 3 and Fig. 1 show the positive significant correlation between bright and black blood techniques. p value = 0.000.

The black blood and the white blood acquisitions were also not significant for the iron loaded ($N = 17$, $p = 0.31$) and non-iron loaded ($N = 33$, $p = 0.081$) subgroups.

The black blood technique was superior to the bright blood technique as regards the study reproducibility ($R^2 1.9 \pm 2.07$ versus 2.4 ± 14.7) in addition to the better inter-observer agreement concerning the black blood technique compared to the bright blood technique (3.2 ± 1.2 versus 8.3 ± 16.4).

4.2. Blood suppression and image quality

Images acquired from midventricular septum using bright- and black-blood sequences (Fig. 2) show the better delineation of myocardial septum borders with less artifacts in black blood T2* images. Blood signal seems contaminating the myocardium in the bright blood image while the myocardium in the black-blood images is homogeneous.

The analysis of the T2* decay curve and the black blood T2* sequence showed an improved curve fit compared to the conventional bright blood sequence (Fig. 3).

In severe iron loaded hearts, the fast signal decay causes the curve to plateau in delayed echo times. Bright blood technique (top image) shows the lower points below the level of background noise (red crosses) causing improper curve fit, so we used the truncation method in the conventional bright blood technique ($R^2 = 0.9983$ with truncation). However, the black blood technique reduced the background noise with better curve fit through the eight gradual increasing echo times, without any truncation ($R^2 = 0.9996$) and this diminishes the analysis errors.

5. Discussion

Iron-overload due to recurrent blood transfusions leading to heart failure related cardiomyopathy is considered the most frequent cause of mortality in patients with thalassemia major. Iron overload is related to repeated blood transfusion, with each unit of blood containing 200–250 mg of elemental iron, along with excessive gastrointestinal adsorption. Transfusional iron is deposited in the reticuloendothelial system (RES); when




Fig. 1 Shows positive significant correlation between bright and black blood techniques.




Fig. 2 (a) Bright blood and (b) black blood techniques of short axis midventricular images obtained from the same thalassemia patient showing the superior image quality and clear septal margins of the black-blood technique.




Fig. 3 Plateau curve in heavily iron loaded heart with rapid decay in the longer echo times. The top image shows the points that fall below background noise level are removed to improve curve fit (red crosses) using the truncation method for the black blood acquisition, and in lower image the background noise is reduced and thus the curve fit is complete for the full range of echo times without truncation which reduces analysis errors.

the stores of the RES are saturated, iron deposition starts to accumulate in parenchymal tissues of the endocrine glands, hepatocytes and myocardium (16,17).

Cardiac MR studies have proved that some thalassemia major patients have a heterogeneous iron deposition pattern in the myocardium as measured by in vivo T2*, although with more and severe cases of iron deposition the heterogeneity begins to diminish (18).

Myocardial T2* in vivo, however, may be confounded by a number of factors, including magnetic susceptibility artifact and measurement error. So in view of these problems, it has been recommended that T2* measurement should be restricted to a full-thickness septal ROI, with highly delineative midseptum R2* measurement (19).

Myocardial T2* and functional CMR imaging, are considered the gold standard for assessment of cardiac iron loading (20). With continuing advances of MRI technology, main field non-uniformity is mainly patient-induced and the T2* is subject to localized susceptibility artifacts (21,22). Fortunately, the susceptibility artifacts can be minimized by detaining the measured ROI to the septum, and subsequently, this technique has demonstrated good reproducibility (23–25) and lately, has been further optimized in terms of trigger delay, echo-spacing, acquisition window and other parameters (26).

A more recent advance was the inclusion of black blood preparation, which provides suppressed blood signal with clearly defined borders, less motion artifacts leading to superior reproducibility and improved interobserver agreement compared to the conventional bright blood GRE sequence (27).

In this study we included 50 TM patients on regular blood transfusion, where 17 had varying degree of cardiac siderosis while 33 were normal as measured with the conventional bright blood and suppression black blood sequences.

Our study objective was to assess the efficacy and reproducibility of black blood IR technique compared to the conventional bright blood GRE sequence in the assessment of cardiac iron concentration.

Our results showed positive significant correlation between the two techniques, with p value 0.000, in agreement with Ou et al. (28), who stated that, the use of a blood suppression prepulse black blood technique had little effect on the calculated cardiac R2*.

We reported superior study reproducibility of black blood T2* imaging in all 50 patients ($R2^*1.9 \pm 2.07$ versus 2.4 ± 14.7) and in the agreement of the inter-observer variability (3.2 ± 1.2 versus 8.3 ± 16.4). Similar results were reported by Smith et al. (29), who found a significant coefficient improvement (three folds) in intra and inter-observer variability with better inter-study reproducibility for black blood T2* imaging compared to the conventional white blood sequence.

Using conventional bright blood sequence, resulting in low signal contrast with difficulty in defining myocardium borders, hence, the blood contamination will be included in the decay curve which increases the possibility of errors in T2* measurement. Misregistration of the high signal blood pool artifacts which can be superimposed on the ROI can also increase the rate of errors.

Using the double inversion recovery black blood technique with better delineation of the myocardial borders, and reducing the noise artifacts will improve the accuracy of T2* measurement (29).

Our results showed better image quality, less artifacts and better delineation of myocardium septum, which improved decay curve analysis and reduced the requirement for truncation method in heavily loaded hearts using the black blood T2* technique, in consistent with Taigang H et al. (30) who believed that the black-blood T2* technique improved potential partial volume errors, reduced artifacts and simplified the data analysis for clinicians and therefore be less sensitive to observer training errors.

6. Conclusion

Black-blood T2* sequence has improved the assessment of myocardial iron concentration, by providing high contrast images, better myocardial border definition, reducing blood signal contamination from the myocardium and less images artifacts. We support the use of blood suppression prepulse for measurement of cardiac T2* based on its superior study reproducibility and interobserver agreement.

Conflict of interest

The authors declare that there are no conflict of interests.

References

- (1) Pennell DJ. T2* magnetic resonance and myocardial iron in thalassemia. *Ann NY Acad Sci* 2005;1054:373–8.
- (2) Borgna-Pignatti C, Rugolotto S, De Stefano P, Zhao H, Cappellini MD, Del Vecchio, et al. Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica* 2004;89:1187–93.
- (3) Modell B, Khan M, Darlison M, Westwood MA, Ingram D, Pennell DJ. Improved survival of thalassaemia major in the UK and relation to T2* cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2008;10:42 (Haematologica 2004;89:1187–93).
- (4) Telfer P. Update on survival in thalassemia major. *Hemoglobin* 2009;33:S76–80.
- (5) Westwood M, Anderson LJ, Firmin DN, Gatehouse PD, Charrier CC, Wonke B, et al. A single breath-hold multiecho T2* cardio vascular magnetic resonance technique for diagnosis of myocardial iron overload. *J Magn Reson Imaging* 2003;18:33–9.
- (6) Olson LJ, Edwards WD, Holmes Jr DR, Miller Jr FA, Nordstrom LA, Baldus WP. Endomyocardial biopsy in hemochromatosis: clinic pathologic correlates in six cases. *J Am Coll Cardiol* 1989;13:116–20.
- (7) Westwood MA, Anderson LJ, Firmin DN, Gatehouse PD, Lorenz CH, Wonke B, et al. Interscanner reproducibility of cardiovascular magnetic resonance T2* measurements of tissue iron in thalassemia. *J Magn Reson Imaging* 2003;18:616–20.
- (8) Kirk P, He T, Anderson LJ, Roughton M, Tanner MA, et al. International reproducibility of single breath hold T2* MR for cardiac and liver iron assessment among five thalassemia centers. *J Magn Reson Imaging* 2010;32:315–9.
- (9) Anderson LJ, Westwood MA, Prescott E, Walker JM, Pennell DJ, Wonke B. Development of thalassaemic iron overload cardiomyopathy despite low liver iron levels and meticulous compliance to desferrioxamine. *Acta Haematol* 2006;115:106–8.
- (10) He T, Gatehouse PD, Kirk P, Tanner MA, Smith GC, Keegan J, et al. Black-blood T2* technique for myocardial iron measurement in thalassemia. *J Magn Reson Imaging* 2007;25:1205–9.
- (11) He T, Gatehouse PD, Kirk P, Mohiaddin RH, Pennell DJ, Firmin DN. Myocardial T2* measurement in iron overloaded tha-

- lassemia: an ex vivo study to investigate optimal methods of quantification. *Magn Reson Med* 2008;60:350–6.
- (12) He T, Gatehouse PD, Kirk P, Tanner MA, Smith GC, Keegan J, et al. Black-blood T2* technique for myocardial iron measurement in thalassemia. *J Magn Reson Imaging* 2007;25:1205–9.
 - (13) Reeder SB, Faranesh AZ, Boxerman JL, McVeigh ER. In vivo measurement of T2* and field inhomogeneity maps in the human heart at 1.5 T. *Magn Reson Med* 1998;39:988–98.
 - (14) Ghugre NR, Enriquez CM, Coates TD, Nelson Jr MD, Wood JC. Improved R2* measurements in myocardial iron overload. *J Magn Reson Imaging* 2006;23:9–16.
 - (15) Pepe A, Positano V, Santarelli MF, Sorrentino F, Cracolici E, De MD, et al. Multislice multiecho T2* cardiovascular magnetic resonance for detection of the heterogeneous distribution of myocardial iron overload. *J Magn Reson Imaging* 2006;23:662–8.
 - (16) Anderson LJ, Westwood MA, Holden S, Davis B, Prescott E, Wonke B, et al. Myocardial iron clearance during reversal of siderotic cardiomyopathy with intravenous desferrioxamine: a prospective study using T2* cardiovascular magnetic resonance. *Br J Haematol* 2004;127:348–55.
 - (17) Pennell DJ, Berdoukas V, Karagiorga M, Ladis V, Piga A, Aessopos A, et al. Randomized controlled trial of deferiprone or deferoxamine in beta-thalassemia major patients with asymptomatic myocardial siderosis. *Blood* 2006;107:3738–44.
 - (18) Positano V, Pepe A, Santarelli MF, Ramazzotti A, et al. Multi slice multi echo T2* cardiac magnetic resonance for the detection of heterogeneous myocardial iron distribution in thalassaemia patients. *NMR Biomed* 2009;22:707–15.
 - (19) Carpenter John-Paul, He Taigang, Kirk Paul, Roughton Michael, et al. On T2* magnetic resonance and cardiac iron. *Circulation* 2011;123(14):1519–28.
 - (20) Wood JC. History and current impact of cardiac magnetic resonance imaging on the management of iron overload. *Circulation* 2009;120:1937–9.
 - (21) Pepe A, Lombardi M, Positano V, Cracolici E, Capra M, Malizia R, et al. Evaluation of the efficacy of oral deferiprone in beta-thalassemia major by multislice multiecho T2*. *Eur J Haematol* 2006;76:183–92 (PubMed: 16451393).
 - (22) Positano V, Pepe A, Santarelli MF, Scattini B, De Marchi D, Ramazzotti A, et al. Standardized T2* map of normal human heart in vivo to correct T2* segmental artefacts. *NMR Biomed* 2007;20:578–90 (PubMed: 17205488).
 - (23) Westwood M, Anderson LJ, Firmin DN, Gatehouse PD, Charrier CC, Wonke B, et al. A single breath-hold multiecho T2* cardiovascular magnetic resonance technique for diagnosis of myocardial iron overload. *J Magn Reson Imaging* 2003;18:33–9 (PubMed: 12815637).
 - (24) Westwood MA, Anderson LJ, Firmin DN, Gatehouse PD, Lorenz CH, Wonke B, et al. Interscanner reproducibility of cardiovascular magnetic resonance T2* measurements of tissue iron in thalassemia. *J Magn Reson Imaging* 2003;18:616–20 (PubMed: 14579406).
 - (25) Tanner MA, He T, Westwood MA, Firmin DN, Pennell DJ. Thalassemia international federation heart T2* investigators. Multi-center validation of the transferability of the magnetic resonance T2* technique for the quantification of tissue iron. *Haematologica* 2006;91:1388–91 (PubMed: 17018390).
 - (26) He T, Tanner MA, Kirk P, Gatehouse PD, Pennell DJ, Firmin DN. Optimization of a multiecho T2* technique for the measurement of myocardial iron overload. *J Cardiovasc Magn Reson* 2006;8:45–7 (abstract 149).
 - (27) He T, Gatehouse PD, Kirk P, Tanner MA, Smith, et al. Black-blood T2* technique for myocardial iron measurement in thalassemia. *J Magn Reson Imaging* 2007;25(6):1205–9 (PubMed: 17520740).
 - (28) Ou Phalla, Zhao Y, Fawal SE, Banka P, Powell AJ. Cardiac T2* measurements in patients with iron overload: a comparison of imaging parameters and analysis techniques. *Magn Reson Imaging* 2012;30:641–8.
 - (29) Smith Gillian C, Carpenter John P, He Taigang, Alamel Mohammed H. Value of black blood T2* cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2011;13:21.
 - (30) He Taigang, Gatehouse Peter D, Kirk Paul, Tanneretal Mark A. Black-blood T2* technique for myocardial iron measurement in thalassemia. *J Magn Reson Imaging* 2007;25:1205–9.